

Two fake Swiss 'towns'

POPULATION SIZE, MIGRATION, DIVERGENCE, ASSIGNMENT, HISTORY

Bayesian inference using the structured coalescent

Migrate-n version 4.2.7 [April-1-2016]

Using Intel AVX (Advanced Vector Extensions)


Compiled for PARALLEL computer architectures

One master and 4 compute nodes are available.

Compiled for a SYMMETRIC multiprocessors (Grandcentral)

Program started at Wed Jul 6 10:57:35 2016

Program finished at Wed Jul 6 11:36:04 2016



Options

Inheritance scalers in use for Thetas:

All loci use an inheritance scaler of 1.0

[The locus with a scaler of 1.0 used as reference]

Random number seed:

(with internal timer) 2131202942

Start parameters:

Theta values were generated

Using a percent value of the prior

M values were generated

Using a percent value of the prior

Connection matrix:

m = average (average over a group of Thetas or M,

s = symmetric migration M, S = symmetric $4Nm$,

0 = zero, and not estimated,

* = migration free to vary, Thetas are on diagonal

d = row population split off column population, D = split and then migration

Population	1	2
1 Ascona	*	0
2 Brissago	d	*

Order of parameters:

1	Θ_1	<displayed>
---	------------	-------------

2	Θ_2	<displayed>
3	$\Delta_{1 \rightarrow 2}$	<displayed>
4	$\sigma_{1 \rightarrow 2}$	<displayed>

Mutation rate among loci:

Mutation rate is constant for all loci

Analysis strategy:

Bayesian inference

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Metropolis sampling

Prior distribution for parameter

Parameter	Prior	Minimum	Mean*	Maximum	Delta	Bins
Theta	Gamma	0.000000	0.010000	0.100000	0.010000	1500
Theta	Gamma	0.000000	0.010000	0.100000	0.010000	1500
M	Gamma	0.000000	500.000000	5000.000000	500.000000	1500
M	Gamma	0.000000	500.000000	5000.000000	500.000000	1500
Splittime mear	Gamma	0.000000	0.100000	1.000000	0.100000	1500
Splittime std	Gamma	0.000000	1.000000	10.000000	1.000000	1500

Markov chain settings:

Long chain

Number of chains

1

Recorded steps [a]

5000

Increment (record every x step [b])

200

Number of concurrent chains (replicates) [c]

10

Visited (sampled) parameter values [a*b*c]

10000000

Number of discard trees per chain (burn-in)

5000

Multiple Markov chains:

Static heating scheme

4 chains with temperatures

1000000.00 3.00 1.50 1.00
Swapping interval is 1

Print options:

Data file:

infile

Haplotyping is turned on:

NO

Output file:

outfile_x0dx

Posterior distribution raw histogram file:

bayesfile

Raw data from the MCMC run:

bayesallfile.gz

Print data:

No

Print genealogies [only some for some data type]:

None

Data summary

Data file: infile

Datatype: Haplotype data

Number of loci: 5

Mutationmodel:

Locus	Sublocus	Mutationmodel	Mutationmodel parameters
-------	----------	---------------	--------------------------

1	1	Felsenstein 84	[Bf:0.24 0.26 0.27 0.22, t/t ratio=2.000]
2	1	Felsenstein 84	[Bf:0.25 0.24 0.26 0.25, t/t ratio=2.000]
3	1	Felsenstein 84	[Bf:0.25 0.24 0.25 0.26, t/t ratio=2.000]
4	1	Felsenstein 84	[Bf:0.26 0.24 0.23 0.27, t/t ratio=2.000]
5	1	Felsenstein 84	[Bf:0.25 0.24 0.27 0.24, t/t ratio=2.000]

Sites per locus

Locus	Sites
1	1000
2	1000
3	1000
4	1000
5	1000

Site rate variation and probabilities:

Locus	Sublocus	Region type	Rate of change	Probability	Patch size
-------	----------	-------------	----------------	-------------	------------

1	1	1	1.000	1.000	1.000
2	1	1	1.000	1.000	1.000
3	1	1	1.000	1.000	1.000
4	1	1	1.000	1.000	1.000
5	1	1	1.000	1.000	1.000

Population	Locus	Gene copies data	(missing)
1 Ascona	1	10	
	2	10	
	3	10	
	4	10	
	5	10	
2 Brissago	1	10	
	2	10	
	3	10	
	4	10	

	5	10	
Total of all populations	1	20	(0)
	2	20	(0)
	3	20	(0)
	4	20	(0)
	5	20	(0)

Bayesian Analysis: Posterior distribution table

Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.00147	0.00473	0.00683	0.00920	0.01547	0.00763	0.00808
1	Θ_2	0.00027	0.00313	0.00503	0.00700	0.01193	0.00563	0.00594
1	$D_{1 \rightarrow 2}$	0.00533	0.00533	0.02700	0.04667	0.04667	0.05833	0.07739
1	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.26000	1.50667	0.26333	0.39478
2	Θ_1	0.00027	0.00313	0.00503	0.00693	0.01180	0.00563	0.00590
2	Θ_2	0.00187	0.00480	0.00683	0.00920	0.01453	0.00763	0.00807
2	$D_{1 \rightarrow 2}$	0.00000	0.00000	0.02633	0.05667	0.22333	0.05700	0.07593
2	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.24667	1.46667	0.25000	0.38263
3	Θ_1	0.00053	0.00340	0.00517	0.00700	0.01140	0.00563	0.00586
3	Θ_2	0.00467	0.00900	0.01203	0.01560	0.02553	0.01350	0.01428
3	$D_{1 \rightarrow 2}$	0.00000	0.00000	0.01900	0.03467	0.13867	0.03500	0.04344
3	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.14667	0.52000	0.15000	0.10798
4	Θ_1	0.00000	0.00207	0.00383	0.00567	0.01080	0.00450	0.00485
4	Θ_2	0.00580	0.01127	0.01383	0.01687	0.02853	0.01543	0.01624
4	$D_{1 \rightarrow 2}$	0.00000	0.01467	0.04033	0.06800	0.20133	0.05767	0.07308
4	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.19333	1.11333	0.19667	0.26516
5	Θ_1	0.00307	0.00453	0.00723	0.01027	0.01253	0.00783	0.00816
5	Θ_2	0.00373	0.00647	0.00817	0.01013	0.01507	0.00957	0.01034
5	$D_{1 \rightarrow 2}$	0.00000	0.00000	0.00633	0.02000	0.05600	0.02033	0.01783
5	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.11333	0.32000	0.11667	0.02610
All	Θ_1	0.00273	0.00473	0.00597	0.00713	0.00933	0.00610	0.00607
All	Θ_2	0.00593	0.00853	0.01010	0.01173	0.01513	0.01037	0.01046
All	$D_{1 \rightarrow 2}$	0.00000	0.00000	0.00033	0.02000	0.05000	0.02033	0.02270
All	$S_{1 \rightarrow 2}$	0.00000	0.00000	0.00333	0.10000	0.28000	0.10333	0.07536

Citation suggestions:





Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters.

Bioinformatics 22:341-345

Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use?

In Population Genetics for Animal Conservation, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli,

and C. Vernesi, eds., vol. 17 of Conservation Biology, Cambridge University Press, Cambridge UK, pp. 42-79.

Bayesian Analysis: Posterior distribution over all loci

Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D | \text{thisModel}) - \ln(\text{Prob}(D | \text{otherModel}))$

or as $LBF = 2(\ln(\text{Prob}(D | \text{thisModel}) - \ln(\text{Prob}(D | \text{otherModel}))$

shows the support for thisModel]

Locus	Raw thermodynamic score(1a)	Bezier approximation score(1b)	Harmonic mean(2)
1	-1926.03	-1806.91	-1800.48
2	-1932.42	-1804.27	-1801.11
3	-2124.71	-1917.93	-1898.24
4	-2508.45	-2185.91	-2146.70
5	-2213.62	-1926.90	-1887.95
All	-10721.58	-9658.29	-9550.84

(1a, 1b and 2) are approximations to the marginal likelihood, make sure that the program run long enough!

(1a, 1b) and (2) should give similar results, in principle.

But (2) is overestimating the likelihood, it is presented for historical reasons and should not be used

(1a, 1b) needs heating with chains that span a temperature range of 1.0 to at least 100,000.

(1b) is using a Bezier-curve to get better approximations for runs with low number of heated chains

[Scaling factor = -16.358429]

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, Genetics, 185: 313-326.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	2089036/6250408	0.33422
Θ_2	2389541/6255166	0.38201
$\Delta_{1 \rightarrow 2}$	3790404/6248321	0.60663
$\sigma_{1 \rightarrow 2}$	1759745/6244215	0.28182
Genealogies	4123414/25001890	0.16492

MCMC-Autocorrelation and Effective MCMC Sample Size

Parameter	Autocorrelation	Effective Sample Size
Θ_1	0.19966	240207.62
Θ_2	0.10919	279042.30
$\Delta_{1 \rightarrow 2}$	0.08952	291212.11
$\sigma_{1 \rightarrow 2}$	0.13242	268512.07
Ln[Prob(D G)]	0.34275	171664.94

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is very common that some parameters for some loci will not be very informative, triggering suggestions (for example to increase the prior range) that are not sensible. This suggestion tool will improve with time, therefore do not blindly follow its suggestions. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. For example, if you run a Bayesian inference with sequence data, for macroscopic species there is rarely the need to increase the prior for Theta beyond 0.1; but if you use microsatellites it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3) it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the range will not help in such situations, reducing number of parameters may help in such situations.

No warning was recorded during the run